



SASSO MANAGEMENT TIPS

MANAGING THE SASSO BREED OF CHICKS



**BIYINZIKA
POULTRY**
INTERNATIONAL LIMITED



SASSO is a dual-purpose bird which is versatile, flexible and easy to manage. It is a coloured bird that is popular amongst consumers and fetches a high price.

Free-range

220 Eggs per annum

70 days, 2 kg

Starts laying at 5 months

COMPARATIVE FEATURES BETWEEN LOCAL CHICKEN AND SASSO

Features	Local chicken	Sasso
Colour	Multi-coloured	Multi-coloured
Egg production per annum	70 Eggs per annum	220 Eggs per annum
Start of eggs	32 weeks	18 weeks
Body weight	1.0 - 1.1kg	2.5 - 5.0kg
Feeding	Household and agriculture waste	Household and agriculture waste
Housing	Rudimentary shelter	Rudimentary shelter
Potential income	Marginal	Substantial

FEEDING SASSO AS A BROILER

FEEDING SASSO AS A LAYER

Feed type	Period	Feed type	Period
Sasso Chick Starter	1 - 18 days	Sasso Chick Starter	1 - 18 days
Broiler Grower	19 - 35 days	Layer Grower Mash	42 - 111 days
Broiler Finisher	36 - sale	Floor Layer Mash	112 - sale

PERFORMANCE OF SASSO AS A BROILER

Age of Bird	Daily Feed Consumption/ Bird	Weekly Body weights	Water Consumption (mls)
7 days	27.3 grams	130 grams	55
14 days	32.8 grams	195 grams	66
21 days	55.4 grams	392 grams	111
28 days	66.5 grams	598 grams	133
35 days	111 grams	893 grams	222
42 days	138.9 grams	1.250 kg	278
49 days	140 grams	1.579 kg	280
56 days	155.3 grams	1.849 kg	311
63 days	169.7 grams	2.286 kg	339
70 days	185 grams	2 kgs	370

VACCINATION SCHEDULE FOR SASSO

PERIOD	TYPE OF DISEASE VACCINATED AGAINST	METHOD OF APPLICATION
Day 1 <input checked="" type="checkbox"/> Done at the hatchery	Mareks (HVT & ND) + Rispens	S/C Injectable
	Newcastle(NCD) + Infectious Bronchitis (IB)	Spray
Day 7	Newcastle(NCD) + Infectious Bronchitis (IB)	Spray/Eye drop
Day 10 - 12	Gumboro	Drinking water
Day 18	Gumboro	Drinking water
Day 21	Newcastle(NCD) + Infectious Bronchitis (IB) Add IB Primer 491	Spray/Eye drop /Drinking water

PERIOD	TYPE OF DISEASE VACCINATED AGAINST	METHOD OF APPLICATION
Week 5	Fowl Pox	Wing stab
Week 10	Fowl Typhoid	Breast muscle
Week 12	Newcastle(NCD) + Infectious Bronchitis (IB)	Spray /Drinking water/Eye drop
Week 14	Infectious Coryza	Subcutaneous
Week 14	Deworming	Drinking water
Week 17	Newcastle(NCD) + Infectious Bronchitis (IB)	Spray /Drinking water/Eye drop
Week 18	Deworming	Drinking water
Every 4 weeks	Newcastle(NCD) + Infectious Bronchitis (IB)	Spray /Drinking water/Eye drop
Every 6 weeks	Add IB Primer/491	Spray

Note that vaccination varies with how long you plan to keep the birds. If you plan to keep the birds for 2 months, use the broiler vaccination schedule.

For technical and sales support contact the area sales managers below:

Kampala-1: 0789 097 422 / 0709 843 349
East: 0761 362 900 / 0702 513 599
Kampala-3: 0786 130 020 / 0752 241 106
North: 0772 384 706 / 0701 849 391

West Mityana, Mubende route:
0788 410 643 / 0703 325 590

Masaka- Mbarara route:
0759 965 052 / 0773 979 795

Biyinzika Poultry International Ltd
Plot 376 Kigunga, Mukono, Uganda.
Tel: +256 712 243 632 / 0800 300 091 (toll free)
Email: info@biyinzika.co.ug